

Applicant : Ernest G. Hope  
Serial No. : 09/722,096  
Filed : November 22, 2000  
Page : 2

Attorney's Docket No.: 12531-002001

A2 cont  
at 10  $\mu$ M BFA. This resistance is sustained after BFA is removed from the media. Resistant lymphoblastoid target AMK were rendered susceptible by a 22 hr TM pre-treatment that was not reversed by BFA treatment. Neither TM or TM and BFA together altered CIK mediated lysis of OCI-Ly8 tumor targets.

---

In the drawings:

Substitute the originally-filed drawings with the formal drawings filed herewith.

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**






FIG. 1




Day 21 CIK effector cell: target cell ratio

FIG. 2A




Day 21 CIK effector cells: OCI-LY8 target ratio 10:1

FIG. 2B




Day 21 CIK effector cell: target cell ratio

FIG. 2C



Day 21 CIK effector cell: target cell ratio 20:1


FIG. 3



Day 21 CIK effector cell: target cell ratio 20:1

FIG. 4





Effector cell: target cell ratio 2:1

FIG. 5




FIG. 6




FIG. 7A





FIG. 7B

# HUVEC Lysis by CIK After Pre-Treatment of HUVEC with Various Concentrations of Purified, Recombinant Human Hsp47



Treatment Group

FIG. 8




FIG. 9






FIG. 10